

Lightest scalars as chiral partners of the Nambu–Goldstone bosons

L. Y. Xiao^a, H. Q. Zheng^{a*} and Z. Y. Zhou^{a†}

^aDepartment of Physics, Peking University, Beijing 100871, China

We review the spectrum of lightest scalar resonances recently determined using dispersion techniques. The conceptual difference between the pole mass and the bare mass (or the line–shape mass) is stressed. The nature of the lightest scalars are discussed and we argue, without relying on any model details, that the $\sigma(500)$, $\kappa(700)$, $a_0(980)$ and $f_0(980)$ may be understood as chiral partners of the Nambu–Goldstone bosons in the linear realization of chiral symmetry. But there remains some difficulties in understanding the role of $f_0(980)$ in this picture.

The question on whether there exists a light and broad resonance with quantum number $IJ^P = 00^+$ below 1GeV (for the historical reason, this resonance is properly called the “ σ ”) had been a long standing issue of debate [1]. The major difficulties to accept σ may be summarized as the following:

1. The σ contribution in $L\sigma M$ must exactly cancel the $\pi\pi$ contact interactions, or more generally, background contributions as dictated by PCAC and the soft pion theorem. In the non-linear realization of chiral symmetry, the sigma meson is hence not needed, at least in the very low energy region.
2. Being light and broad makes the σ ’s contribution to the phase shift hard to be distinguishable from background contributions while the latter are often out of control in various phenomenological studies.
3. The $\pi\pi$ phase shift in the $I=0$, $J=0$ channel does not pass $\pi/2$ at moderately low energies which should have appeared if there exist a light σ as required by the standard Breit–Wigner description of resonances.

The above questions can be answered. In elastic $\pi\pi$ scatterings, the background contribution, or the left hand cut contribution can be estimated at low energies using chiral perturbation theory.

*Invited talk given by Zheng at QCD06, Montpellier, France, July 3 – 7, 2006.

†Present address: Department of Physics, Southeast University, Nanjing 210096, China

It is found that the background contribution to the $IJ=00$ channel $\pi\pi$ scattering phase shift is negative and concave while the experimental data is positive and convex, hence demonstrating that the σ meson is crucial to adjust chiral perturbation theory to experiments [2], and hence the existence of σ is a must. Furthermore, in PKU parametrization form [3,4,5], which may be considered as an upgraded Dalitz–Tuan parametrization form, partial wave elastic scattering amplitudes are constructed using dispersion technique, and the amplitudes satisfy analyticity and single channel unitarity by only assuming Mandelstam analyticity. In PKU form the pole contribution to the phase shift is constructed explicitly and it is easy to verify that, when there appears a large width, the pole mass parameter, denoted as m , can be totally different from the mass scale M where the phase shift contributed by the pole pass $\pi/2$ [3]. Taking the σ resonance for example, while $m \sim 450\text{MeV}$, M is of order 1GeV. The two mass parameters m and M looks quite different, but are actually only different faces of a single light and broad resonance. The PKU form nicely depicts and explains the strong enhancement of $IJ=00$ channel $\pi\pi$ scattering phase shift over a long range of the center of mass energy, which is named as “red dragon” by Minkowski and Ochs [6]. Fig. 1 depicts the situation, where one sees clearly how a large width distorts the pole contribution to the phase shift, comparing with the typical narrow resonance $f_0(980)$ ’s contribution. The background contribution is not depicted in fig. 1, since its contribution is found




Figure 1. The σ pole contribution to the $IJ=00$ channel $\pi\pi$ scattering phase shift. See Ref. [4] for more information.

to be small numerically. According to this, it is possible to define the mass parameter M for a pole properly. Actually in the PKU parametrization form each pole's contribution to the S matrix is factorized:

$$S^{pole}(s) = \frac{M^2(z_0) - s + i\rho(s)sG[z_0]}{M^2(z_0) - s - i\rho(s)sG[z_0]}, \quad (1)$$

where

$$M^2(z_0) = \text{Re}[z_0] + \frac{\text{Im}[z_0] \text{Im}[z_0 \rho(z_0)]}{\text{Re}[z_0 \rho(z_0)]}, \quad (2)$$

$$G[z_0] = \frac{\text{Im}[z_0]}{\text{Re}[z_0 \rho(z_0)]}.$$

In above the mass parameter M is given as a function of the pole mass ($\sqrt{z_0} \equiv m + i\Gamma/2$), ρ is the kinematic factor. The pole mass parameters of σ and κ are determined to be (in units of MeV) [4,5]:

$$\begin{aligned} m_\sigma &= 470 \pm 50, & \Gamma_\sigma &= 570 \pm 50, \\ m_\kappa &= 694 \pm 53, & \Gamma_\kappa &= 606 \pm 59. \end{aligned} \quad (3)$$

These results, though with larger error bars, are fully compatible with the excellent results obtained using Roy equations: $m_\sigma = 441_{-8}^{+16}$ MeV,

$\Gamma_\sigma = 544_{-8}^{+25}$ MeV [7] and $m_\kappa = 658 \pm 13$ MeV, $\Gamma_\kappa = 557 \pm 24$ MeV [8]. Comparing with the sophisticated Roy equation analyses which demonstrate the existence of the sigma and kappa by direct calculation, the PKU approach is more intuitive, simpler, and without much loss of rigorousness (the theoretical uncertainties from the left hand cut in the latter approach is highly suppressed). It is worth emphasizing that, even if there is no Mandelstam representation, and hence analyticity of partial wave amplitudes is not established on the whole complex plane, it is proved over a large region [9] which is much larger than the analyticity domain straightforwardly established from Roy equations, and is certainly large enough for any phenomenological application. The PKU parametrization form is remarkable in the sense that it is sensitive to the isolated singularities not too far from the elastic threshold. Except the determination on σ and κ , it is also found to be useful in correctly figuring out the virtual state pole in the $I=2$ s wave exotic channel [4], as well as the tiny effect of a virtual pole from the $I=0$ d wave $\pi\pi$ scattering phase shift data [10]. Comparing with the Roy equation analysis, the one adopted in the present approach are orthogonal calculations, and it is important to have calculations using different methods on such important outputs and that the final results agree with each other. The value on the pole positions of σ and κ obtained using dispersion techniques, provide useful constraints when discussing the structure of the lightest QCD scalars.

The first important information is the large width of σ and κ . The large widths are quite often overlooked when discussing the mass spectrum in the literature. It is often attempted to set up SU(3) mass relations among pole mass parameters m . However, a light σ with a mass around 500 MeV as a bare parameter appeared in the lagrangian can hardly produce a large width, in any model calculations. On the other side, the parameter M for σ or κ can be determined using Eq. (2) and the estimated value of the pole positions are $M_\sigma \simeq 930$ MeV and $M_\kappa \simeq 1380$ MeV with sizable error bars. It is noticed that the parameter M is very much like the bare mass parameter appeared in a simple K matrix approach, or the

renormalized mass in the propagator of loop calculations, since all of them take the role of determining where the phase shift contributed by the resonating field passes $\pi/2$. Parameter M may also be called as the line-shape mass. Qualitatively speaking the stronger the resonance couples to the $\pi\pi$ continuum, the larger the deviation is between m and M . The situation is illustrated in fig. 2. If the comparison is correct, it will lead to




Figure 2. The pole trajectory determined from Eq. (1) as an illustrative example. The bare mass is taken as 1GeV. The pole mass decreases as the coupling strength, G , increases. The unit is MeV.

an important conceptual change when discussing the possible SU(3) mass relations among lightest scalars: σ , κ , $a_0(980)$ and also $f_0(980)$. Instead of comparing different m , one should first examine the relations between different “bare” masses, M . Since the former quantities associated with large widths are severely distorted by the strong couplings to the pseudo-goldstone pairs, it is not suitable to use them to discuss the SU(3) mass relations. For example, we have

$$m_\sigma < m_\kappa < m_{a_0} , \quad (4)$$

but actually the mass relation should be read as

$$1\text{GeV} \simeq M_\sigma \leq M_{a_0} < M_\kappa . \quad (5)$$

We will argue in the following that the relation Eq. (5) is consistent with a chiral $\bar{q}q$ octet description to the lightest scalars. Especially the bare mass of κ exceeds M_{a_0} , hence the original motivation leads to the tetra quark description of the lightest scalars [11] already disappear. In fact it can be proved under mild assumptions, that a tetra quark state (or more precisely, a tetra quark as the leading component in Fock state expansions in the large N_c limit) does not exist since it violates analyticity [12]. There exist a few mass relations among the light scalars taking into account 't Hooft's instanton effects, in the extended Nambu–Jona-Lasinio model [13]:

$$M_{f_0}^2 + M_{f_0'}^2 \simeq 2M_\kappa^2 - g_A M_{tH}^2 , \quad (6)$$

$$M_\kappa^2 - M_{a_0}^2 \simeq 3g_A(M_K^2 - M_\pi^2) , \quad (7)$$

where $M_{tH}^2 = M_\eta^2 + M_{\eta'}^2 - 2M_K^2 \simeq 0.72\text{GeV}^2$ characterizes the QCD anomaly contribution to the scalar sector, and g_A is the axial vector coupling due to $\pi - A_1$ mixing. The above formulas are obtained by assuming that the scalars are chiral partners of the pseudo-goldstone bosons. They are not exact quantities but may serve as a crude “first order” approximation. From the mass relations it is found that a smaller value of M_κ and g_A is preferred. Taking $g_A \simeq 0.6$ for example [14], if taking $M_\kappa = 1.5\text{GeV}$, then through Eq. (7) one would attempt to consider the $a_0(1450)$ as a member of the chiral scalar octet. But then the well established conventional $\bar{q}q$ state $K_0^*(1430)$ will miss its own SU(3) partner, which is not permitted. If taking $M_\kappa = 1.2\text{GeV}$, then Eq. (7) tells $M_{a_0} \simeq 1.02\text{GeV}$, i.e., not far from $a_0(980)$. The major difficulty in this picture comes from the mass of f_0 . Eq. (6) indicates $M_{f_0} \simeq 1.2$, if taking $M_{f_0'=\sigma} \simeq 1\text{GeV}$. Taking into account the couple channel effects, which are large for $f_0(980)$, does not seem to be helpful in resolving the discrepancy. Consider a simple Flatté formula for a couple-channel Breit–Wigner propagator,

$$BW(s) \propto \frac{1}{M^2 - s - i\rho_1 g_1^2 - i\rho_2 g_2^2} . \quad (8)$$

where the subscripts denotes different channels. For a large M^2 above the second threshold this expression generates a broad third sheet resonance and may also generate a light resonance

below the $\bar{K}K$ threshold. But unfortunately the light one can only be located on the 4th sheet whereas the observed $f_0(980)$ is on the second sheet. For $f_0(980)$ the situation is more complicated since its bare state f_0 can also mix with $f_0(1370)$, etc. Without instanton effects, the σ and f_0 are ideally mixed and the latter is $|\bar{s}s\rangle$. When the instanton effects are taken into account f_0 also contains sizable $|\bar{n}n\rangle$ content and hence may have a sizable mixing with $f_0(1370)$, thus reducing significantly its mass. Hence we think that the bare mass of $f_0(980)$ being roughly 250MeV above the observed state is not a true difficulty in accepting the picture that the σ , κ , $a_0(980)$ and $f_0(980)$ comprise the chiral partners of the pseudo-goldstone octet. This idea is not new [15]. The above discussions may however be helpful in resolving some confusions and misconceptions widely spread in the literature. One simple way to understand the difference between the pole mass m and the bare mass M is through a proper unitarization procedure [16]. One advantage of the chiral model with linearly realized chiral symmetry is that it is not difficult to explain at qualitative level the large width of σ and especially the large width of κ . It is more difficult to explain the latter [17], unless it possesses a bare mass much larger than the pole mass parameter. At lower energies one can then further integrate out the scalar fields as well as other light resonances to get a lagrangian with non-linearly realized chiral symmetry. We stress that the “linear σ model” here implies a more realistic linear chiral model rather than the BPHZ renormalizable toy σ model, since the latter failed to reproduce the LECs of the Gasser–Leutwyler lagrangian [18].

The use of dispersion techniques firmly establishes the existence of σ and κ , and also provides a reliable determination to their mass parameters. The physical meaning of these mass parameters are clarified and it is proposed that the lightest scalars are chiral partners of the pseudo-goldstone bosons. Nevertheless the role of $f_0(980)$ in such picture remains to be clarified.

Acknowledgement: One of the author, H.Z., would like to thank the organizer of QCD06, Stephan Narison, for providing a charming atmosphere during the conference and helpful dis-

cussions, and also Wolfgang Ochs, for interesting and helpful discussions in the related subject. Especially H.Z. is grateful to Prof. San Fu Tuan, for his kind encouragement. The authors would also like to thank Zhi-Hui Guo and Ming-Xian Su for helpful discussions. This work is supported in part by National Natural Science Foundation of China under contract number 10575002 and 10421503.

REFERENCES

1. For a review, see N. A. Tornqvist, hep-ph/0008135.
2. Z. G. Xiao and H. Q. Zheng, Nucl. Phys. **A695** (2001) 273; Chin. Phys. Lett. **20** (2003) 342.
3. H. Q. Zheng *et al.*, Nucl. Phys. **A733** (2004) 235.
4. Z. Y. Zhou *et al.*, JHEP0502 (2005) 43.
5. Z. Y. Zhou and H. Q. Zheng, Nucl. Phys. **A755** (2006) 212 (hep-ph/0603062 v2).
6. P. Minkowski, W. Ochs, hep-ph/9905250.
7. I. Caprini, G. Colangelo and H. Leutwyler, Phys. Rev. Lett. **96** (2006) 132001.
8. S. Descotes-Genon, B. Moussallam, hep-ph/0607133.
9. A. Martin, Nuovo Cim. **44** (1966) 1219.
10. J. J. Wang, Z. Y. Zhou and H. Q. Zheng, JHEP 0512 (2005) 019.
11. R. L. Jaffe, Phys. Rev. **D15** (1977) 267.
12. Z. G. Xiao and H. Q. Zheng, hep-ph/0502199, to appear in MPLA.
13. M. X. Su *et al.*, to appear. It is in qualitative agreement with A. A. Osipov, H. Hansen, B. Hiller, Nucl. Phys. **A745**(2004)81.
14. J. Bijnens, C. Bruno and E. de Rafael, Nucl. Phys. **B390**(1993)501.
15. See for example, N. Törnqvist, Euro. Phys. J. **C11**(1999)359.
16. See for example, D. Black *et al.*, Phys. Rev. **D64** (2001) 014031.
17. This is seen in QSSR approach, see, S. Narison, hep-ph/0512256, and also private communications.
18. J. Gasser and H. Leutwyler, Annals of Physics, **158** (1984) 142.